arXiv:1203.0854v1 [math.DG] 5 Mar 2012

A NOTE ON BLOW-UPS OF TORIC SURFACES AND CSC KÄHLER METRICS

CARL TIPLER

Abstract. Let X be a compact toric surface. Then there exists a sequence of torus equivariant blow-ups of X such that the blown-up toric surface obtained admits a cscK metric.

1. Introduction

The problem of finding constant scalar curvature Kähler metrics (cscK for short) on compact Kähler manifolds is of main interest in Kähler geometry since the work of Yau [23], Tian [21] and Donaldson [5]. The Tian-Yau-Donaldson conjecture asserts that there exists a cscK metric on a polarized Kähler manifold (X, L) in the Kähler class $c_1(L)$ if and only if the pair (X, L) is stable in a suitable GIT sense. At that time, we know that the existence of a cscK metric implies the K-polystability by a result of Stoppa [19]. Stoppa's argument relies on an important result of Arezzo and Pacard on blow-ups of cscK manifolds [1], [2]. Arezzo and Pacard managed to show that if (X, ω) is cscK, then the blow-up of X at finitely many well chosen points admits a cscK metric. In order to prove their result, they use a gluing method. They build the blown-up manifold by gluing a local model, the blow-up of $\mathbb{C}^n$ at 0, endowed with the Burns-Simanca metric [18], which is ALE and scalar flat.

Another result on blow-ups and special metrics is the one of Taubes [20] that asserts that there are anti-self-dual metrics on the blow-up at sufficiently many points of any smooth oriented 4-manifold. In the Kähler case, a metric is anti-self-dual if and only if it is scalar-flat.

This two results suggest the following:

Conjecture 1. *Let X be a compact Kähler manifold. Then there exists a finite sequence of blow-ups of X such that the blown-up manifold obtained admits a cscK metric.*

In the case of ruled surfaces, this conjecture has been proved by Le Brun and Singer [11] for surfaces of genus higher than 2 and then in general by Kim, Le Brun and Pontecorvo [10]. However, in their argument, if the starting surface is toric, the resulting blown-up surface does not admit a torus symmetry anymore.

On the other hand, Rollin and Singer found another method to built cscK metrics on blow-ups of surfaces, based on parabolic structures [16], from which we can deduce a lot of examples of toric surfaces with cscK metrics.

In this paper, we show that if the initial surface is toric, then we can prove the conjecture staying in the toric world.

Theorem 1. *Let X be a compact complex smooth toric surface. Then there exists a finite sequence of blow-ups of X at torus fixed points such that the blown-up toric surface obtained admits a cscK metric.*

Remark 1. *By a remark of Wang and Zhou* [22]*, using the theorem of Arezzo, Pacard and Singer on blow-ups of extremal metrics* [3]*, we know that each toric surface admits an extremal metric. However from Matsushima and Lichnerowicz obstruction, we know from Calabi* [4] *that the Hirzebruch surfaces admit no extremal metric of constant scalar curvature. Thus we need to blow-up toric surfaces if we want them to admit cscK metrics.*

Remark 2. *About the existence of Kähler-Einstein metrics on toric manifolds, a recent result of Legendre* [12] *shows that each toric orbifold admits a unique singular Kähler-Einstein metric.*

The proof of this result relies on simple considerations on the classification of toric surfaces and the result of Arezzo and Pacard on blow-ups of cscK metrics. We hope that it will ouline a stabilization process of blowing-up a manifold at suitable points and its link with the existence of cscK metrics. Basically, the idea is to blow-up the toric surface until it has a good shape. We first build a sequence of toric surfaces :

$$... \to X_{j+1} \to X_j \to ... \to X_1 \to X_0$$

with $X_0 = \mathbb{P}^1 \times \mathbb{P}^1$ and X_{j+1} obtained from X_j by blowing-up each torus fixed-point. By the theorem of Arezzo and Pacard, each of these X_j admits a cscK metric.

Then, using the classification of toric surfaces, we prove that for each smooth toric surface X, there exists a sequence of blow-ups at torus fixed points such that the blown-up surface obtained is one of the X_j. From the fan descrition point of view, we add rays to the fan describing X untill it becomes very symmetric, isomorphic to the fan of one of the X_j. In fact it turns out that the X_j endowed with a suitable polarization are K-polystable and asymptotically Chow polystable.

We will deduce from our result the following corollary:

Corollary 2. *Let X be a smooth compact toric surface. Then there is a sequence of toric blow-ups of X (see section 2):*

$$Bl_{(p_m,..,p_1)} \to X$$

and a polarization $L \to Bl_{(p_m,..,p_1)}$ such that $(Bl_{(p_m,..,p_1)}, L)$ is K-polystable and asymptotically Chow-polystable.

Remark 3. *Donaldson has proved that a polarized toric surface (X, L) admits a cscK metric in the class $c_1(L)$ if and only if (X, L) is K-polystable* [6]*. However it is a hard problem, even for toric surfaces, to check K-polystability. It contrasts with our construction which is explicit. We can compute from the fan describing the toric surface the points to blow-up in order to reach one of the X_j.*

Corollary 2 lies on deep results of Mabuchi [13] and Stoppa [19] and computation of the Lie algebra characters associated to Mabuchi's obstruction. This computation is simplified by the symmetries of the surfaces.

Remark 4. *The idea of using symmetries on toric polytopes has also been used by Donaldson* [5]*, but in order to construct a 9 points blow-up of $\mathbb{CP}^2$ that is not K-polystable for the associated polarization, even if its Futaki invariant vanishes.*

Thus this polarized toric surface admits no cscK metric in the corresponding Kähler class.

We will also compute a bound on the number of blow-ups necessary to get a cscK metric with our construction. We will see by an example that this bound is far from being optimal, raising the natural question:

Question 3. *What is the minimal number of blow-ups necessary to obtain a cscK metric on an iterated blow-up of the n^{th} Hirzebruch surface $\mathbb{F}_n$?*

We point out that we know from Wang and Zhou [22] that each toric surface admits an extremal metric. This metric is cscK if and only if the Futaki invariant vanishes [8] and an explicit formula for this invariant depending on the polytope is known since the work of Donaldson [5]. However this is far from answering the question as the Kähler classes of the extremal metrics and the kähler classes where the Futaki invariant vanishes might be far from each other.

1.1. Plan of the paper.

In section 2 we recall results on blow-ups of toric surfaces. Then the proof of theorem 1 is given in section 3. The last section deals with stability issues related to our construction.

1.2. **Acknowledgements.** I would like to thank professors Vestislav Apostolov, Paul Gauduchon, Yann Rollin and Michael Singer for their constant support.

2. Blow-ups of toric CSC surfaces

2.1. Toric surfaces.

A toric surface X is a complex algebraic surface endowed with an effective action of a complex torus $T^{\mathbb{C}}$ of dimension 2, such that there exists an open and dense orbit in X. Thus X is a compactification of $T^{\mathbb{C}}$ and is obtained by gluing divisors to this torus. The structure of X is then entirely described by the way these divisors are glued together, and this procedure is encoded by a fan $\Sigma = \Sigma(X)$ in a $\mathbb{Z}$-lattice N, [14]. A fan Σ is the union of strongly convex rationnal polyhedral cones σ in $N_{\mathbb{R}} = N \otimes_{\mathbb{Z}} \mathbb{R}$ satisfying two gluing conditions:

- $\forall \sigma \in \Sigma$, $\forall \tau$ face of σ, $\tau \in \Sigma$
- $\forall (\sigma_1, \sigma_2) \in \Sigma^2$, $\sigma_1 \cap \sigma_2$ is a face of σ_1 and σ_2

To each cone σ in Σ, one can associate an open affine toric variety $U_\sigma = \mathrm{spec}(\mathbb{C}[\sigma^{>} \cap M])$, where M is the dual lattice of N and

$$\sigma^{>} = \{R \in M / \langle R, x\rangle \geq 0 \ \forall x \in \sigma\}.$$

Then the gluing conditions of the fan ensure that this affine charts glue together and form a toric variety X.

This gives a simple description of these surfaces as each of their algebraic properties admits a combinatorial conterpart in term of the fan.

We will denote by $\Sigma^{(i)}(X)$ the set of cones of the fan describing X of dimension equal to i. We will identify the rays in $\Sigma^{(1)}$ and primitive generators of these rays.

Example 2.1.1. Let $\mathbb{F}_n$ be the n^{th} Hirzebruch surface, that is the total space of the fibration

$$\mathbb{P}(\mathcal{O} \oplus \mathcal{O}(n)) \to \mathbb{CP}^1.$$

Note that $\mathbb{F}_0 = \mathbb{CP}^1 \times \mathbb{CP}^1$. Then $\Sigma^{(1)}(\mathbb{F}_n) = \{(0,-1),(1,0),(0,1),(-1,-n)\}$ in $N = \mathbb{Z}^2$. As an example $\mathbb{F}_2$ admits the fan description of figure 1.



Figure 1. $\mathbb{F}_2$

Here we only represent the elements of $\Sigma^{(1)}(\mathbb{F}_2)$ by arrows. The other cones of Σ are $\{0\}$ and the two dimensional cones delimited by the rays of $\Sigma^{(1)}$. We will use this way of describing fans in the remaining of this paper.

2.2. **Blow-ups.** The blow-up of a toric manifold $X(\Sigma)$ along a torus invariant divisor is described in term of the fan Σ. To each cone $\sigma \in \Sigma$, one can associate a torus invariant divisor V_σ as the closure of spec$(\mathbb{C}[\sigma^\perp \cap M])$ in X, where $\sigma^\perp = \{R \in M / \langle R, x\rangle = 0 \forall x \in \sigma\}$. Then if σ is of full dimension, V_σ is a torus fixed-point. Let τ be the ray in $N_\mathbb{R}$ generated by the sum of the primitive generators of dimension one faces of σ. Then τ induces a subdivision of σ into rationnal strictly convex polyhedral cones, $\sigma = \cup_i \sigma_i$. Then the blow-up of X at V_σ is a toric manifold described by the fan $\Sigma - \{\sigma\} \cup_i \{\sigma_i\}$.

Example 2.2.1.
Consider the cone $\sigma_1 \in \Sigma(\mathbb{F}_2)$ generated by $(0,1)$ and $(-1,-n)$ in the lattice $N = \mathbb{Z}^2$. The blow-up of $\mathbb{F}_2$ at the point $p = V(\sigma)$ is pictured on figure 2.



Figure 2. $Bl_p\mathbb{F}_2$

We will say that a blow-up is *toric* if it is the blow-up at torus fixed points, so that the action of the torus lifts to the blown-up manifold. We introduce the following notation: if X is the blow-up of Y at a point $p \in Y$, we set $X = Bl_p(Y)$. Then by proposition 2.2.2, as $\mathbb{F}_1 \simeq Bl_p(\mathbb{CP}^2)$ for p a torus fixed point, any toric surface X different from $\mathbb{CP}^2$ can be written $Bl_{p_m}(Bl_{p_{m-1}}(...(Bl_{p_1}(\mathbb{F}_n))))$ for some $n \geq 0$ and $p_{j+1} \in Bl_{p_j}(Bl_{p_{j-1}}(...(Bl_{p_1}(\mathbb{F}_n))))$. We will use the notation $Bl_{(p_m,..,p_1)}(\mathbb{F}_n) = Bl_{p_m}(Bl_{p_{m-1}}(...(Bl_{p_1}(\mathbb{F}_n))))$ and call it an *iterated blow-up*.

The combinatorial description of singular points and completness for toric surfaces lead to the following result [7]:

Proposition 2.2.2. *Each smooth compact toric surface is obtained from an iterated toric blow-up of* $\mathbb{CP}^2$ *or one of the Hirzebruch surfaces* $\mathbb{F}_n$.

2.3. The Arezzo and Pacard theorem.

In the proof of theorem 1, we will build a sequence of toric surfaces, each admitting a cscK metric. This sequence is built by successive blow-ups. We recall the following theorem of Arezzo and Pacard [1] on blow-ups of cscK manifolds:

Theorem 2.3.1. *Let* (X, ω) *be a cscK compact Kähler manifold and let* $(p_j)_{j=1..m}$ *be* m *distinct points of* X. *Assume that* X *admits no holomorphic vector field. Consider the blow-up of* X *at the* p_j:

$$\pi : Bl_{(p_j)}(X) \to X.$$

Then for each $a_j \in \mathbb{R}^{+*}$ *positive weights, there exists* $\varepsilon_0 > 0$ *such that for each* $\varepsilon \in (0, \varepsilon_0)$, *there is a cscK metric on* $Bl_{(p_j)}(X)$ *in the Kähler class*

$$\pi^*[\omega] - \varepsilon^2 \sum_j a_j PD(E_j)$$

where $PD(E_j)$ *denotes the Poincaré dual of the exceptionnal divisor* $E_j = \pi^{-1}(p_j)$.

In a second paper, Arezzo and Pacard proved a similar result when X admits holomorphic vector fields. However in that case the blow-up points need to satisfy some stability and genericity conditions [2]. An other way to use their theorem in that case is to perform the analysis modulo a finite group of automorphisms that preserves no holomorphic vector field:

Theorem 2.3.2. *Let* (X, ω) *be a cscK compact Kähler manifold. Let* $G \subset \mathrm{aut}(X)$ *be a finite group of isometries of* ω *and* $G \cdot p_1 = \{p_1, .., p_m\}$ *be the orbit of a point* $p_1 \in X$. *Let* r_j *be the order of the stabilizer of* p_j *in* G. *Assume moreover that no holomorphic vector field is* G*-invariant. Consider the blow-up of* X *at the* p_j:

$$\pi : Bl_{(p_j)}(X) \to X.$$

Then for each $a \in \mathbb{R}^{+*}$, *there exists* $\varepsilon_0 > 0$ *such that for each* $\varepsilon \in (0, \varepsilon_0)$, *there is a cscK metric on* $Bl_{(p_j)}(X)$ *in the Kähler class*

$$\pi^*[\omega] - \varepsilon^2 a \sum_j r_j PD(E_j)$$

where $PD(E_j)$ *denotes the Poincaré dual of the exceptionnal divisor* $E_j = \pi^{-1}(p_j)$.

3. Proof of the main theorem

Let X be a compact toric surface. We want to show that a torus equivariant blow-up of X admits a cscK metric. We will first build a sequence of toric surfaces X_j endowed with cscK metrics and then show that we can reach one of these X_j by an iterated blow-up of X.

3.1. The surfaces X_j.

The X_j are built inductively. Let $X_0 = \mathbb{CP}^1 \times \mathbb{CP}^1$. Then X_{j+1} is the blow-up of X_j at its 2^{j+2} torus fixed points. Using the fan description of the toric blow-up process, the fan describing $\mathbb{CP}^1 \times \mathbb{CP}^1$, X_1 and X_2 are represented on figure 3, 4 and 5.



FIGURE 3. $\mathbb{CP}^1 \times \mathbb{CP}^1$



FIGURE 4. X_1



FIGURE 5. X_2

Proposition 3.1.1. *Each of the X_j admits a cscK metric.*

Proof. We proceed by induction. First $\mathbb{CP}^1 \times \mathbb{CP}^1$ admits a cscK product metric. Then we apply the Arezzo and Pacard theorem 2.3.2. Consider the group G in $\mathrm{aut}(X_0)$ generated by the following elements, described in terms of their action in homogeneous coordinates:

$$([u_1, v_1], [u_2, v_2]) \longmapsto ([v_1, u_1], [u_2, v_2]),$$

$$([u_1, v_1], [u_2, v_2]) \longmapsto ([u_1, v_1], [v_2, u_2]),$$

$$([u_1, v_1], [u_2, v_2]) \longmapsto ([-u_1, v_1], [u_2, v_2])$$

and

$$([u_1, v_1], [u_2, v_2]) \longmapsto ([u_1, v_1], [-u_2, v_2]).$$

Then the G orbit of $([1,0],[1,0])$ consists of four fixed points under the torus action induced by the $\mathbb{C}^*$ action on each $\mathbb{CP}^1$. No holomorphic vector field of X_0 is invariant under the G-action thus by theorem 2.3.2 X_1 admits a cscK metric. Note that the weights of the exceptionnal divisors in X_1 are the same.

By induction, we build from a cscK metric on X_j, $j \geq 1$, a cscK metric on X_{j+1}. Note that for each $j \geq 1$, the Lie algebra of $\mathrm{aut}(X_j)$ is the Lie algebra of the complex two torus that defines the toric structure. Note also that the group in $\mathrm{aut}(X_0)$ generated by

$$([u_1,v_1],[u_2,v_2]) \longmapsto ([v_1,u_1],[u_2,v_2])$$

and

$$([u_1,v_1],[u_2,v_2]) \longmapsto ([u_1,v_1],[v_2,u_2])$$

lifts to a group G_j in $\mathrm{aut}(X_j)$. The holomorphic vector fields generating the torus action are not invariant under the action of G_j. We now iterate Arezzo and Pacard theorem, working modulo G_j at each step and blowing-up the orbits of each torus fixed points, giving the same weight to each new exceptionnal divisor. Note that by construction of the cscK metrics, the group G_j acts by isometries at each step and we can iterate the blow-up process. □

3.2. **From X to X_j.**

To end the proof of the theorem we will use the two following lemmas:

Lemma 3.2.1. *For any sequence of m toric blow-ups of $\mathbb{F}_n$, $Bl_{(p_m,..,p_1)}(\mathbb{F}_n)$, one of the following holds:*

- *$n = 0$ or*
- *there is a sequence of m toric blow-ups of $\mathbb{F}_{n-1}$ such that $Bl_{(p_m,..,p_1)}(\mathbb{F}_n) = Bl_{(q_m,..,q_1)}(\mathbb{F}_{n-1})$ or*
- *there is a point $p_{m+1} \in Bl_{(p_m,..,p_1)}(\mathbb{F}_n)$ such that there is a sequence of $m+1$ toric blow-ups of $\mathbb{F}_{n-1}$ such that $Bl_{(p_{m+1},..,p_1)}(\mathbb{F}_n) = Bl_{(q_{m+1},..,q_1)}(\mathbb{F}_{n-1})$*

and

Lemma 3.2.2. *For any iterated toric blow-up of X_0, $Y = Bl_{(q_m,..,q_1)}(X_0)$, there is an iterated toric blow-up of Y such that*

$$Bl_{(q'_r,..,q'_1)}(Y) = X_j$$

for some j.

The classification of toric surfaces is the key for the proof of the theorem:

Proof of Theorem 1. By proposition 2.2.2, there exists a sequence of toric blow-ups such that $X = Bl_{(p_m,..,p_1)}(\mathbb{F}_n)$.

By iterating lemma 3.2.1, there are toric blow-ups $Bl_{(q_m,..,q_1)}(X)$ of X such that $Bl_{(q_m,..,q_1)}(X)$ is an iterated blow-up of $\mathbb{F}_0 = X_0$.

Then by lemma 3.2.2 we can blow-up $Bl_{(q_m,..,q_1)}(X) = Bl_{(q'_r,..,q'_1)}(X_0)$ to reach one of the X_j. By proposition 3.1.1, this blow-up of X has a cscK metric. □

We end this section by the proof of the two lemmas:

Proof of lemma 3.2.1. Suppose $n \neq 0$. Consider the fan describing $\mathbb{F}_n$ on figure 6.

Then a blow-up of $\mathbb{F}_n$ is described by adding a ray generated by the sum of two adjacent primitive vectors in $\Sigma^{(1)}(\mathbb{F}_n)$ to this fan. Each further blow-up is described by the same process so we can separate the blow-ups of $\mathbb{F}_n$ into two types, those



FIGURE 6. $\mathbb{F}_n$

arising by adding a ray in the cone $\sigma_1 = \mathbb{R}^+(0,1) + \mathbb{R}^+(-1,-n)$, called type one blow-ups, and the other. These two types of blow-ups commutes, we can perform every blow-up of type 1 before the other blow-ups to obtain $Bl_{(p_m,..,p_1)}(\mathbb{F}_n)$. Then there are two possibilities. If there is no blow-up of type 1, we blow-up V_{σ_1} and we are in the third case of lemma 3.2.1. Indeed, the blow-up of $\mathbb{F}_n$ at V_{σ_1} is a one-point blow-up of $\mathbb{F}_{n-1}$, see figure 7.



FIGURE 7. $Bl_p(\mathbb{F}_n) = Bl_q(\mathbb{F}_{n-1})$

In the other case, the first blow-up of type one is necessarily at V_{σ_1} and we are in the second case of lemma 3.2.2. $\square$

We need to introduce some definitions. We say that an iterated toric blow-up $Bl_{(p_m,..,p_1)}(X)$ is a *purely iterated blow-up* if for each $i \geq 2$, p_i lies on the exceptional divisor E_{i-1} coming from the blow-up at p_{i-1}. In that case we call m the *length* of this iterated blow-up. Let X be a toric surface and n be the minimal integer such that X is obtained from toric blow-ups of $\mathbb{F}_n$. Then we define the integer $l(X)$ to be the maximal length of purely iterated blow-ups arising in the description of X from $\mathbb{F}_n$.

Proof of lemma 3.2.2. Let $\Sigma^{(1)}(Y)$ be the set of rays of the fan of Y, identified with primitive generators of these rays. Then the fan description of a toric blow-up implies that there is a sequence of fans Σ_j such that $\Sigma_0 = \Sigma(X_0)$, $\Sigma_m = \Sigma(Y)$ and $\Sigma^{(1)}_{j+1}$ is obtained from $\Sigma^{(1)}_j$ by adding a ray generated by $v_i + v_{i+1}$, with v_i and v_{i+1} two adjacent elements of $\Sigma^{(1)}_j$. Consider the set of all $m+1$-tuples $(v_0, .., v_m)$ of $\Sigma^{(1)}_m$ such that $v_i \in \Sigma^{(1)}_i$, and for each $i \geq 1$, $v_i = v_{i-1} + w_{i-1}$ with $w_{i-1} \in \Sigma^{(1)}_{i-1}$ adjacent to v_{i-1} or $w_{i-1} = 0$. The case when $w \neq 0$ correspond to the blow-ups. Denote $l(v_0, .., v_m)$ the number of distinct elements in $(v_0, .., v_m)$ and let $l_0(Y)$ be the maximum of the $l(v_0, .., v_d)$. Then $l_0(Y) = l(Y) + 1$. Then we can blow-up Y in order to make its fan symmetrical untill we obtain the fan describing $X_{l(Y)}$. It is enough to add the rays that are missing so that we can get every $m+1$-tuple $(v_0, .., v_m)$ with $l_0(v_0, .., v_m) = l_0(Y)$ which are coming from the set of rays of $X_{l(Y)}$. An example is pictured on figure 8.

Example 3.2.3.



Figure 8. Y

On this example $d = 2$ because of the sequence $\{(0,1),(1,1),(1,2)\}$. We blow-up 5 times to add symmetries, as represented on figure 9.



Figure 9. adding symmetries

And we blow-up again to reach X_2. □

3.3. **Bound on the number of blow-ups.** We prove the following upper bound on the minimal number of blow-ups that are needed to get a cscK metric with our method:

Proposition 3.3.1. *Let X be a toric surface. Let n be the minimal integer such that X is obtained from toric blow-ups of $\mathbb{F}_n$. Then there is a sequence of at most*

$$2^{n+l(X)+1} - \sharp(\Sigma^{(1)}(X))$$

blow-ups of X such that the blown-up surface admits a cscK metric.

Remark 3.3.2. Note that for $X = \mathbb{F}^n$ we have $l(X) = 1$ and $2^{n+2} - 4$ is the number of blow-ups necessary to go from $\mathbb{F}_n$ to X_n and in that case this estimate is sharp for our method.

Proof. We need to estimate the number of blow-ups necessary to go from X to the nearest X_j. As X is a blow-up of $\mathbb{F}_n$ for n minimal, there is a n times blow-ups of X such that $Bl_{(q_n,..,q_1)}(X)$ is a blow-up of $X_0 = \mathbb{F}_0$. Then from the proof of lemma 3.2.2, we know that there exists an iterated blow-up from $Bl_{(q_n,..,q_1)}(X)$ to $X_{n+l(X)-1}$. So we can go from X to $X_{n+l(X)-1}$ by an iterated blow-up. Then the number of blow-ups necessary is bounded by the number of rays in $\Sigma^{(1)}(X_{n+l(X)-1})$ minus the number of rays in $\Sigma^{(1)}(X)$, which gives the result. $\square$

This result on the minimal number of points to be blown-up in order to get a cscK metric is not sharp. Indeed, from [17], we know that the special blow-up of $\mathbb{F}_2$ pictured on figure 10 admits a cscK metric.



FIGURE 10. A special blow-up of $\mathbb{F}_2$

Here we do not need to blow-up further in order to reach X_2. It suggests the following problem:

Question 3.3.3. *What is the minimal number of blow-ups of $\mathbb{F}_n$ necessary to ensure the existence of a cscK metric on the blown-up surface?*

4. Blown-up surfaces and stabilities

In GIT several notions of stability for polarized varieties are studied. It is in general a hard problem to check that a polarized variety is stable or not. However, the existence of a cscK metric enables to conclude for K-stability and asymptotic Chow stability in some cases. In this section, we deduce from theorem 1 that each toric surface can be blown-up to satisfy K-stability and asymptotic Chow stability for some polarization.

4.1. **X_j and K-stability.** We refer the reader to [19] for the definition of K-stability. We can assume that the cscK metrics ω_j built on the X_j lie in rationnal classes, so that up to scaling, we can assume that these metrics represent a polarization L_j of X_j, that is $[\omega_j] = c_1(L_j)$. By a result of Stoppa [19], the (X_j, L_j) are K-polystable. Together with theorem 1 we have:

Corollary 4.1.1. *Let X be a smooth compact toric surface. Then there is a sequence of toric blow-ups of X:*

$$Bl_{(p_m,..,p_1)} \to X$$

and a polarization $L \to Bl_{(p_m,..,p_1)}$ such that $(Bl_{(p_m,..,p_1)}, L)$ is K-polystable.

4.2. **Asymptotic Chow-polystability of the X_j.** We refer to [9] for the definition of asymptotic Chow-stability. By a theorem of Mabuchi [13], if (X, L) is a polarized Kähler manifold with cscK metric in the class $c_1(L)$ and if Mabuchi's obstruction vanishes, then (X, L) is asymptotically Chow polystable. From the work of Ono [15] and Futaki [9], Mabuchi's obstruction admits a simple description for toric varieties. Assume that (X, L) is a polarized toric manifold of complex dimension n, with L a torus equivariant line bundle. Suppose that the lattice $N = \mathbb{Z}^n$. Then this polarization defines a polytope $\Delta \subset \mathbb{R}^n$ that encodes the symplectic structure of X [14]. Then set for each $i \in \{0, .., n\}$

$$\mathbf{s}_\Delta(i) = \sum_{\mathbf{a} \in i\Delta \cap \mathbb{Z}^n} \mathbf{a}$$

and

$$E_\Delta(i) = \sharp(i\Delta \cap \mathbb{Z}^n).$$

Then

$$i \mapsto \mathrm{Vol}(\Delta)\mathbf{s}_\Delta(i) - iE_\Delta(i) \int_\Delta \mathbf{x} dv = \sum_j i^j \mathcal{F}_{\Delta,i}$$

is a polynomial in i with coefficients $\mathcal{F}_{\Delta,i}$, with dv the euclidian volume of $\mathbb{R}^n$. Then Futaki has shown that the vanishing of Mabuchi's obstruction is equivalent to the vanishing of the $\mathcal{F}_{\Delta,i}$.

Lemma 4.2.1. *Let Δ_j be the Delzant polytope associated to the toric polarized surface (X_j, L_j). Then $\mathcal{F}_{\Delta_j,1} = \mathcal{F}_{\Delta_j,2} = 0$.*

Proof. This is a straightforward computation, using the symmetries of Δ_j with respect to the two coordinate axes. Note that $\mathcal{F}_{\Delta,1}$ is the Futaki invariant and thus necessarily vanishes by the existence of the cscK metric. □

From this lemma and Mabuchi's theorem we deduce:

Corollary 4.2.2. *Let X be a smooth compact toric surface. Then there is a sequence of toric blow-ups of X:*

$$Bl_{(p_m,..,p_1)} \to X$$

and a polarization $L \to Bl_{(p_m,..,p_1)}$ such that $(Bl_{(p_m,..,p_1)}, L)$ is asymptotically Chow-polystable.

References

[1] C.Arezzo and F.Pacard, *Blowing up and desingularising constant scalar curvature Kähler manifolds*, Acta Math. **196** (2006), no. 2, 179-228.

[2] C.Arezzo and F.Pacard, *Blowing up constant scalar curvature Kähler manifolds II*, Ann. of Math. (2) **170** (2009), no. 2, 685-738.

[3] C.Arezzo, F.Pacard and M.Singer, *Extremal Metrics on blow ups.* Duke Math. J. Volume **157**, Number 1 (2011), 1-51.

[4] E. Calabi, *Extremal Kähler metrics*, Seminars on Differential Geometry (S. T. Yau Ed.), Annals of Mathematics Studies, Princeton University Press, 1982, pp. 259-290.

[5] S.K.Donaldson, *Scalar curvature and stability of toric varieties.* J. Differential geometry **62**, (2002), 289-349.

[6] S.K.Donaldson, *Constant scalar curvature metrics on toric surfaces.* Geom. Func. Anal., **19** (2009), 83-136.

[7] W.Fulton. *Toric Varieties.* Princeton University Press 1993.

[8] A.Futaki *Kähler-Einstein metrics and Integral Invariants*, Springer-LNM **1314**, Springer-Verlag (1988).

[9] A.Futaki, *Asymptotic Chow polystability in Kähler geometry*, preprint ArXiv 1105.4773v2.

[10] J.-S. Kim, C.Le Brun and M. Pontecorvo *Scalar-Flat Kähler Surfaces of All Genera*, J. reine angew. Math. **486** (1997) 69-95.

[11] C.Le Brun and Michael Singer, *Existence and Deformation Theory for Scalar-Flat Kähler Metrics on Compact Complex Surfaces*, Inv. Math. **112** (1993) 273-313.

[12] E.Legendre, *Toric Kähler-Einstein metrics and convex compact polytopes*, preprint ArXiv 1112.3239.

[13] T.Mabuchi, *An energy-theoretic approach to the Hitchin-Kobayashi correspondance for manifolds, I*, Invent. Math. **159** (2005), 225-243.

[14] T. Oda, *Convex bodies and algebraic geometry*, Ergebnisse der Mathematik und ihrer Grenzgebiete, 3. Folge, Band 15, Springer-Verlag, (1988).

[15] H.Ono, *A necessary condition for Chow semistability of polarized toric manifolds*, to appear in J.Math.Soc.Japan.

[16] Y.Rollin and M.Singer, *Constant scalar curvature Kähler surfaces and parabolic polystability.* J. Geom. Anal. **19** (2009), no. 1, 107-136.

[17] Y.Rollin, C.Tipler, *Deformations of extremal toric manifolds*, preprint ArXiv 1201.4137.

[18] S.R. Simanca, *Kähler metrics of constant scalar curvature on bundles over* $\mathbb{CP}_{n-1}$, Math. Annalen **291**, (1991), no.2, 239-246.

[19] J.Stoppa, *K-stability of constant scalar curvature Kähler manifolds*, Adv. Math, **221** (2009), 1397-1408.

[20] C.H.Taubes, *The existence of Anti-Self-Dual metrics*, J.Diff. Geometry, **36**, (1992), 163-253.

[21] G.Tian, *Kähler-Einstein metrics with positive scalar curvature*, Invent. Math. **137**, (1997), 1-37.

[22] Xu-jia Wang, Bin Zhou, *On the existence and nonexistence of extremal metrics on toric Kähler surfaces*, Advances in Mathematics **226**, (2011), 4429-4455.

[23] S.-T. Yau, *Open problems in geometry*, Proc. Symposia Pure Math. **54**, (1993), 1-28.

Département de Mathématiques, Laboratoire Jean Leray, 2, Rue de la Houssinière - BP 92208, F-44322 Nantes, FRANCE

E-mail addresses: `carl.tipler@univ-nantes.fr`